

Glossary For Prosthetists

EDITOR'S NOTE: *This brief glossary was obtained from Mr. Steven Purka, Clerk in the New York City office of the Prosthetic and Sensory Aids Service, Veterans Administration. The Glossary was compiled from a number of sources. Readers will recall that Mr. Purka was the compiler of "Good Reading for Growing Technicians", which appeared in the JOURNAL of OALMA, August, 1951, 33-36.*

GENERAL

Anatomy—The science of the *structure* of the body and relation of its parts.

Biomechanics—The science which deals with *forces* which act on the body.

Kinesiology—The study of muscular and joint *movements*.

Physiology—The science of the *functions* of the body and its parts.

Rehabilitation—The restoration to useful activity of individuals who have been injured so as to suffer from physical or emotional disability, such restoration including treatment of the disability and training to fit the individual for occupation in industry.

Therapy—The science that deals with *diseases and remedies*. *Occupational therapy* is the use of an occupation or training in an occupation for remedial purposes. *Physical therapy* is the treatment of disease by physical means.

ANATOMICAL

Abduction—The withdrawal of a part from the axis of the body; of the foot—rotation of the foot outward, on its own axis. To move away or to be away from the mid-line of the body. (For a part to be further away from the mid-line than normal).

Adduction—Any movement whereby a part is brought toward another or toward the median line of the body. A part of the body is nearer the mid-line of the body than normal when in adduction.

Acromion—The lateral projection of the spine of the scapula forming the point of the shoulder.

Anterior—Situated in front of or in the forward part of.

Arm—The portion of the upper extremity between the acromion and the epicondyles.

Axilla—The armpit.

Cervical Vertebra—7th—The most prominent of all the vertebrae. When the head is bent forward, the 7th cervical vertebra is the prominent bulge at the back of the neck.

Clavicle—The collar bone.

Condyle—A prominence or swelling of a bone at a joint, especially when occurring in pairs. Condyles of the tibia ordinarily take bearing in a prosthesis for amputation below the knee.

Distal—Remote, farthest from the center, origin, or point of attachment. (Opposite: Proximal).

Dorsal—Pertaining to the back or to the posterior part of an organ or of the body.

Epicondyles—The bony prominences of the distal end of the humerus.

Eversion—A turning outward. Eversion of the foot, turning the sole away from the mid-line of the body.

Extension—A straightening out, especially the muscular movement by which a flexed limb is made straight.

Femur—The thigh bone.

Fibula—The outer, smaller bone of the leg below the knee.

Flexion—The act of bending; the condition of being bent.

Forearm—The portion of the upper extremity between the epicondyles and the styloids.

Humerus—The bone of the upper part of the arm.

Hyperextension—Excessive extension. Hyperextension of the knee is that beyond 180° as by stretching of the hamstrings in the normal knee or by compression of a bumper in an artificial knee.

Inferior—Lower. Also toward the feet.

* Reviewed in the Veterans Administration and published with the approval of the Chief Medical Director. The statements and conclusions published by the author are the result of his own study and do not necessarily reflect the opinion or policy of the Veterans Administration.

Inferior Angle of the Scapula—The angle of the lower point of the scapula.

Inversion—The act of turning inward.

Ischium—The lower and rearmost of the principal bones in either half of the pelvis; the seat bone. In ischial bearing, the weight of the body is supported on the rear wall of the top of the prosthesis as on a bicycle seat.

Lateral—Toward the outside. (Opposite: Medial).

Medial—Toward the inside or center. (Opposite: Lateral).

Neuroma—The new growth made up of nerve cells and nerve fibers found at the end of cut nerves.

Normal Limb—A remaining, unamputated limb.

Palmar—Pertaining to the palm of the hand.

Patella—The knee cap.

Plantar—Pertaining to the sole of the foot.

Posterior—Placed behind or to the back of a part; behind.

Pronation—The act of turning the palm of the hand downward. (Opposite: Supination).

Proximal—Nearest to the body, or center, or some other point considered as the center of a system.

Radius—The shorter, thicker bone of the forearm.

Residual—Remaining in the stump, refers to motion, function, etc.

Rotation—Movement about an axis.

Sagittal—Pertaining to the anteroposterior median plane of the body (sagittal plane), the median vertical plane of the body dividing it into right and left halves.

Styloid of the Ulna—The bony prominence at the lateral side of the wrist when the palm is turned down.

Superior—Higher, denoting the upper of two parts. Also towards the head.

Supination—The turning of the palm of the hand upward. (Opposite: Pronation).

Tibia—The inner, front, larger bone of the leg below the knee.

Transverse—Crosswise; at right angles to the longitudinal axis of the body.

Trochanter—Bony prominence at the top of the femur.

Ulna—The inner, longer bone of the forearm.

ABBREVIATIONS

AE—Above-elbow.

AK—Above-knee.

BE—Below-elbow.

BK—Below-knee.

CKJ—Effective center of rotation of knee joint, about $\frac{3}{4}$ " above the joint space.

EB—End-bearing. Also Elbow-bearing. (Disarticulation by analogy).

KB—Knee-bearing. (Disarticulation.)

OPERATIONS

Chopart—An amputation of the foot consisting of a disarticulation through the tarsal joints, leaving only the os calcis and the astragalus.

Cineplasty—A plastic surgical procedure to produce a muscle motor used to initiate motion of a prosthesis.

Disarticulation—Separation at a joint; amputation through a joint—the proximal part of the joint being left in the stump.

Gritti-Stokes—(For amputation above the knee-joint). The patella is preserved in a long anterior flap, and, having had a thin slice removed from its deep surface, is secured in apposition with the femur, the latter having been deprived of its articular surface by being sewn through the condyles. Stokes—the same as Gritti's operation, except that section of the femur is made above the condyles. Intended to give end-bearing.

Lisfranc—A disarticulation of the metatarsal bones from the tarsus.

Pirogoff—(For amputation through the foot). A partial osteoplastic operation in which the os calcis is sawed through obliquely from above and downward and forward, and the posterior portion is brought up and secured against the surface made by sawing off the lower ends of the tibia and fibula. Intended to give end-bearing stump; formerly used in Europe.

Syme's—Amputation at the ankle joint, the malleoli being sawn through and a flap made with the skin of the heel. Intended to give end-bearing.

Tenoplasty—Plastic surgery of a tendon.

PREFERRED USAGE

Suction Socket—(Preferred to vacuum socket). Socket for limb held to the stump by wearer with atmospheric pressure when the limb is unsupported.

Socket—(Preferred to "bucket"). Any container for amputation stump; may or may not be integral with structural member.

